

Black Holes on the Brane with Induced Gravity

G. Kofinas

*Department of Physics, University of Crete,
710 03 Heraklion, Crete, Greece*

E. Papantonopoulos and V. Zamarias

*Department of Physics, National Technical University of Athens,
Zografou Campus GR 157 80, Athens, Greece*

September 30, 2002

Abstract. An analysis of a spherically symmetric braneworld configuration is performed when the intrinsic curvature scalar is included in the bulk action. In the case when the electric part of the Weyl tensor is zero, all the exterior solutions are found; one of them is of the Schwarzschild- $(A)dS_4$ form, which is matched to a modified Oppenheimer-Volkoff interior solution. In the case when the electric part of the Weyl tensor is non zero, the exterior Schwarzschild- $(A)dS_4$ black hole solution is modified receiving corrections from the non-local bulk effects. A non-universal gravitational constant arises, depending on the density of the considered object and the Newton's law is modified for small and large distances; however, the conventional limits are easily obtained.

Keywords: Black Holes, Branes

1. Introduction

Branes are solitonic solutions of ten-dimensional string theories. In the most simplified picture of the braneworld scenario, our physical world is realized as a four-dimensional hypersurface embedded in a five-dimensional space called bulk. All matter and gauge interactions live on the brane, while the gravitational interactions are effective in the whole five-dimensional space. This allowed to define a gravitational scale of the whole space, which if the extra dimensions are large, can be as low as the TeV scale (Antoniadis et al., 1998 ; Randall and Sundrum, 1999), while the four-dimensional gravitational scale of our world is at the Planck scale.

The effective brane equations have been obtained (Shiromizu, Maeda and Sasaki, 2000) when the effective low-energy theory in the bulk is higher-dimensional gravity. If the dynamics is governed not only by the ordinary five-dimensional Einstein-Hilbert action, but also by the four-dimensional Ricci scalar term induced on the brane, new phenomena appear. It was observed (Dvali, Gabadadze and Porati, 2000), that the localized matter fields on the brane (which couple to bulk gravitons) can generate via quantum loops a localized four-dimensional worldvolume



© 2008 Kluwer Academic Publishers. Printed in the Netherlands.

kinetic term for gravitons. That is to say, four-dimensional gravity is induced from the bulk gravity to the brane worldvolume by the matter fields confined to the brane.

We will discuss static spherically symmetric solutions of braneworlds with induced gravity (Kofinas, Papantonopoulos and Pappa, 2001), and we will present an exterior Schwarzschild- $(A)dS_4$ solution which is matched to a modified interior Oppenheimer-Volkoff solution. In this solution the gravitational constant get corrected for very small matter densities. The conventional solar system bounds of General Relativity set the crossover scale below the TeV scale. The above results were obtained by setting the electric part of the Weyl tensor, $E_{\mu\nu}$ to zero.

Then, we will generalize our study by including the non-local bulk effects (Kofinas, Papantonopoulos and Zamarias, 2002), as they are expressed by a non-vanishing electric part of the Weyl tensor on the brane. By choosing $g_{tt} = -g_{rr}^{-1}$, the system of equations on the brane becomes closed and all the possible static black hole solutions are found for these metrics. These solutions have generic new terms which give extra attractive force compared to the Newtonian $-(A)dS_4$ force, and represent the strong-gravity corrections to the Schwarzschild- $(A)dS_4$ spacetime.

2. Four-Dimensional Static Spherically Symmetric Solutions

We consider a 3-dimensional brane Σ embedded in a 5-dimensional spacetime. Capital Latin letters $A, B, \dots = 0, 1, \dots, 4$ will denote full spacetime, lower Greek $\mu, \nu, \dots = 0, 1, \dots, 3$ run over brane worldvolume. For convenience, we can quite generally, choose a coordinate y such that the hypersurface $y = 0$ coincides with the brane. The total action for the system is taken to be:

$$S = \frac{1}{2\kappa_5^2} \int \sqrt{-^{(5)}g} (^{(5)}R - 2\Lambda_5) d^5x + \frac{1}{2\kappa_4^2} \int \sqrt{-^{(4)}g} (^{(4)}R - 2\Lambda_4) d^4x \\ + \int \sqrt{-^{(5)}g} L_5^{mat} d^5x + \int \sqrt{-^{(4)}g} L_4^{mat} d^4x. \quad (1)$$

We basically concern on the case with no fields in the bulk, i.e. $^{(5)}T_{AB} = 0$. From the dimensionful constants κ_i^2 , the Planck masses M_i are defined as $\kappa_i^2 = 8\pi G_{(i)} = M_i^{i-2}$. Then, a distance scale r_c is defined as $r_c = M_4^2/M_5^3$.

Varying (1) with respect to the bulk metric g_{AB} , and reducing the resulting equations to four dimensions (Kofinas, 2001), we get four-dimensional Einstein gravity, coupled to a well-defined modified matter

content. More explicitly, one gets

$${}^{(4)}G_\nu^\mu = \kappa_4^2 {}^{(4)}T_\nu^\mu - \left(\Lambda_4 + \frac{3}{2}\alpha^2\right) \delta_\nu^\mu + \alpha \left(L_\nu^\mu + \frac{L}{2} \delta_\nu^\mu\right), \quad (2)$$

where $\alpha \equiv 2/r_c$, while the quantities L_ν^μ are related to the matter content of the theory through the equation

$$L_\lambda^\mu L_\nu^\lambda - \frac{L^2}{4} \delta_\nu^\mu = \mathcal{T}_\nu^\mu - \frac{1}{4} (3\alpha^2 + 2\mathcal{T}_\lambda^\lambda) \delta_\nu^\mu, \quad (3)$$

and $L \equiv L_\mu^\mu$. The quantities $\mathcal{T}_\nu^\mu$ are given by the expression

$$\begin{aligned} \mathcal{T}_\nu^\mu = & \left(\Lambda_4 - \frac{1}{2}\Lambda_5\right) \delta_\nu^\mu - \kappa_4^2 {}^{(4)}T_\nu^\mu + \\ & + \frac{2}{3} \kappa_5^2 \left({}^{(5)}\overline{T}_\nu^\mu + \left({}^{(5)}\overline{T}_y^y - \frac{{}^{(5)}\overline{T}}{4} \right) \delta_\nu^\mu \right) - \overline{\mathbf{E}}_\nu^\mu. \end{aligned} \quad (4)$$

Bars over ${}^{(5)}T_\nu^\mu$ and the electric part $\mathbf{E}_\nu^\mu$ of the Weyl tensor mean that the quantities are evaluated at $y = 0$. The presence of $\overline{\mathbf{E}}_\nu^\mu$ makes the brane equations (2) not to be, in general, closed (Shiromizu, Maeda and Sasaki, 2000). Due to the contracted Bianchi identities, the following differential equations among L_ν^μ arise from (2) $L_{\nu;\mu}^\mu + L_{;\nu}/2 = 0$. We are looking for solutions of (2) for spherically symmetric braneworld metrics

$$ds_{(4)}^2 = -B(r)dt^2 + A(r)dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2). \quad (5)$$

The matter content of the 3-universe is considered to be a localized spherically symmetric perfect fluid. We first consider the case $\overline{\mathbf{E}}_\nu^\mu = 0$ as the boundary condition of the propagation equations in the bulk space. All the solutions outside a static localized matter distribution were found. One of these is the Schwarzschild- $(A)dS_4$ metric which is matched to a modified Oppenheimer-Volkoff interior. The exterior solution was found to be

$$B_{>}(r) = \frac{1}{A_{>}(r)} = 1 - \frac{\gamma}{r} - \beta r^2, \quad r \geq R, \quad (6)$$

where γ is an integration constant and

$$\beta = \frac{1}{3}\Lambda_4 + \frac{1}{2}\alpha^2 - \frac{\alpha}{2\sqrt{3}} \sqrt{4\Lambda_4 - 2\Lambda_5 + 3\alpha^2}. \quad (7)$$

Considering a uniform distribution $\rho(r) = \rho_o = \frac{3M}{4\pi R^3}$ for the object, the interior solution was found, and its matching to the above exterior specified the integration constant γ . The result is

$$\frac{1}{A_{<}(r)} = 1 - \left(\beta + \frac{\gamma}{R^3}\right) r^2, \quad r \leq R, \quad (8)$$

$$B_{<}(r) = \frac{1 - \frac{\gamma}{R} - \beta R^2}{\left(1 + \frac{4\pi R^3}{3M} p(r)\right)^2}, \quad r \leq R, \quad (9)$$

$$p(r) = -\rho_o \frac{\sqrt{1 - (\beta + \frac{\gamma}{R^3})r^2} - \sqrt{1 - (\beta + \frac{\gamma}{R^3})R^2}}{\sqrt{1 - (\beta + \frac{\gamma}{R^3})r^2} - \omega \sqrt{1 - (\beta + \frac{\gamma}{R^3})R^2}}, \quad (10)$$

where

$$\gamma = \frac{\kappa_4^2 M}{4\pi} + \frac{\alpha R^3}{2\sqrt{3}} \sqrt{4\Lambda_4 - 2\Lambda_5 + 3\alpha^2} - \frac{\alpha R^3}{2\sqrt{3}} \sqrt{4\Lambda_4 - 2\Lambda_5 + 3\alpha^2 + \frac{3\kappa_4^2 M}{\pi R^3}}, \quad (11)$$

$$\omega^{-1} = 1 - \frac{2}{\kappa_4^2 \rho_o} \left(\beta + \frac{\gamma}{R^3} \right) \left(1 - \frac{\sqrt{3}\alpha}{\sqrt{4\Lambda_4 - 2\Lambda_5 + 3\alpha^2 + 4\kappa_4^2 \rho_o}} \right)^{-1}. \quad (12)$$

The parameters γ and β of the Schwarzschild-(A)dS₄ exterior solution (6) can be constrained by solar system experiments. The bounds obtained fix the crossover scale below the *TeV* range. The γ parameter in the $1/r$ term modifies the Newton's gravitational constant which, as it is seen from (11), and for small matter it densities deviates significantly from its conventional value. This is shown in Fig. 1.




Figure 1. The ρ_0 dependence of Newton's constant in various models: (1) The Newton's constant variation in our model; If $4\Lambda_4 - 2\Lambda_5 \gg 3\alpha^2$ then (1a), otherwise (1b). (2) The behaviour of Newton's constant in another class of our solutions. (3) The Newton's constant variation in RS model.

So far we considered local corrections to the Einstein equations on the brane. The presence of the electric part $\bar{E}^\mu_\nu$ of the Weyl tensor in (4) indicates the 5D gravitational stresses, which are known as massive KK modes of the graviton and this term cannot be ignored.

The symmetric and traceless tensor $\bar{\mathbf{E}}_{\mu\nu}$ is uniquely and irreducibly decomposed as follows

$$\bar{\mathbf{E}}_{\mu\nu} = \mathcal{U}\left(u_\mu u_\nu + \frac{1}{3}\mathbf{h}_{\mu\nu}\right) + \mathcal{P}_{\mu\nu} + 2\mathcal{Q}_{(\mu}u_{\nu)}, \quad (13)$$

where $\mathbf{h}_{\mu\nu} = g_{\mu\nu} + u_\mu u_\nu$ is the projection operator normal to u^μ , while $\mathcal{U}$ is the non-local energy density, $\mathcal{P}_{\mu\nu}$ the non-local anisotropic stress, and $\mathcal{Q}_\mu$ the non-local energy flux on the brane. Static spherical symmetry implies (Maartens, 2002) that

$$\mathcal{Q}_\mu = 0 \quad , \quad \mathcal{P}_{\mu\nu} = \mathcal{P}(r)\left(r_\mu r_\nu - \frac{1}{3}\mathbf{h}_{\mu\nu}\right), \quad (14)$$

where r_μ is the unit radial vector. We will discuss here one class of solutions of (2) with non-zero $\mathbf{E}_{\mu\nu}$. This is

$$\begin{aligned} B = \frac{1}{A} &= 1 - \frac{\gamma}{r} - \beta r^2 \\ &+ sg(\zeta) \frac{\delta}{r} \left[\frac{128}{105} {}_1F_1\left(\frac{15}{8}, \frac{23}{8}; sg(\zeta)z\right) z \right. \\ &\left. + \frac{9}{8} \left(\frac{1}{z} - sg(\zeta)\frac{8}{7}\right) e^{sg(\zeta)z} \right] z^{\frac{7}{8}} \end{aligned} \quad (15)$$

where δ and γ are integration constants (γ is typically interpreted as $2G_N M$ with M being the mass of the point particle, G_N the Newton's constant),

$$\beta = \frac{1}{3}\Lambda_4 + \frac{1}{2}\alpha^2 \quad , \quad \zeta = \frac{\alpha^2}{9}(4\Lambda_4 - 2\Lambda_5 + 3\alpha^2), \quad (16)$$

and $r = \left(\frac{\delta}{\sqrt{|\zeta|}}\right)^{\frac{1}{3}} z^{\frac{1}{8}} e^{sg(\zeta)z/3}$. The electric components of the Weyl tensor are given by the equation, $\mathcal{P} = -2\mathcal{U} = 9\zeta/(2\alpha^2)$. Comparing the solution (15) with the solution (6) having $\bar{\mathbf{E}}_\nu^\mu = 0$, we note that there is a new term, besides the conventional Newtonian and $(A)dS_4$ terms, which carries the information of the gravitational field in the bulk. This new term modifies the Newton's law for small and large distances as it is shown in Fig. 2.

3. Conclusions

We presented new black hole solutions on the brane with induced gravity. Solving the full theory taking under consideration the non-local bulk effects, we found static spherically symmetric solutions, which




Figure 2. Dotted lines represent the Newtonian force. Continuous lines represent the total force, i.e. the sum of the Newtonian and the new force.

represent strong gravity corrections to the Schwarzschild- $(A)dS_4$ black hole solutions. Their characteristic is that they predict a new attractive force. There are classes of solutions with increasing r , where this attractive force combined with the Newtonian one, results to a net force which decreases slower than the Newton's force. In another class of solutions with decreasing r , the new force starts to deviate from the Newton's force at small distances, indicating that at submillimeter scale we could have testable deviations from the Newtonian law. We have also found modifications to Newton's law as a result of a change of the Newton's constant due to the finite interior of the rigid object.

References

- I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. B436 (1998), hep-ph/9804398; N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Rev. D59 (1999) 086004, hep-th/9807344.
- L. Randall and R. Sundrum, Phys. Rev. Lett. 83 (1999) 3370, hep-th/9905221; Phys. Rev. Lett. 83 (1999) 4690, hep-th/9906064.
- T. Shiromizu, K. Maeda and M. Sasaki, Phys. Rev. D62 (2000) 024012, gr-qc/9910076.
- G. Dvali, G. Gabadadze and M. Porati, Phys. Lett. B485 (2000) 208, hep-th/0005016; G. Dvali and G. Gabadadze, Phys. Rev. D63, 065007 (2001), hep-th/0008054.
- G. Kofinas, E. Papantonopoulos and I. Pappa, to appear in Phys. Rev. D, hep-th/0112019.
- G. Kofinas, E. Papantonopoulos and V. Zamarias, to appear in Phys. Rev. D, hep-th/0208207
- G. Kofinas, JHEP 0108 (2001) 034, hep-th/0108013.
- R. Maartens, Phys. Rev. D62 (2000) 084023, hep-th/0004166; gr-qc/0101059.